



Aruna Seneviratne MD
 Tel. 212-636-8290
 Fax 212-636-3102

Department of Orthopaedic Surgery
 425 West 59th Street, 5th floor
 New York, NY 10019

BICEP TENODESIS REHABILITATION PROTOCOL

	RANGE OF MOTION	SLING	EXERCISES
PHASE I 0-4 weeks	PROM as tolerated*	0-4 weeks: except for exercises and hygiene	0-2 weeks: wrist/hand ROM, pendulums, wall stretch at home per instruction sheets 2-4 weeks: Grip strengthening, pulleys/canes, wrist/hand ROM, pendulums; closed chain scapula Deltoid cuff isometrics *Avoid active biceps until 6 wks Begin scapular protraction/retraction
PHASE II 4-8 weeks	Increase as tolerated to full	No Sling	Advance isometrics with arm at side, rotator cuff and deltoid Advance to therabands and dumbbells as tolerated, capsular stretching at end-ROM to maintain flexibility *Avoid active biceps until 6 wks
PHASE III 8-12 weeks	Progress to full without discomfort	No Sling	Advance strength training as tolerated Begin eccentrically resisted motions and closed chain activities Advance to sport and full activity as tolerated after 12 weeks

*If a distal clavicle excision is performed, horizontal adduction should be minimized until pain resolves